



4R Liquid Manure Management Strategies for Optimizing Right Time, Right Place

by Christine Brown, CCA-ON, 4R-NMS

Manure storage capacity will always be an issue if opportunities for application are not considered until the storage is full. To avoid potential issues, look for manure application opportunities during the year and particularly during the growing season and within the crop rotation. The chart below provides an overview of manure application options that will check all the boxes to meet good 4R nutrient stewardship practices.

Month		Application Opportunities/Precautions with Liquid Manure
All Year	<ul style="list-style-type: none"> ✓ ✓ ✓ X 	<ul style="list-style-type: none"> • Sample manure at time of application to analyze nutrient contribution for crops • Use analysis information to fine-tune crop needs from commercial fertilizer • Calibrate application equipment to ensure proper application rate/distribution • Avoid applying manure on fields with very high soil fertility levels
March	✓	<ul style="list-style-type: none"> • Consider building additional storage where current storage is inadequate (approvals and building permits take time)
April	✓	<ul style="list-style-type: none"> • Consider covering an existing storage to reduce the cost of transporting water • Manure applied to winter wheat (monitor: topography and liquid loading) • Manure applied to crop residue or ahead of spring tillage • Injection or surface application followed by immediate incorporation maximizes soil/manure contact
May	✓	
June	<ul style="list-style-type: none"> X X X 	
June	<ul style="list-style-type: none"> ✓ ✓ 	<ul style="list-style-type: none"> • Maximize application to growing crops (↑nutrient efficiency; ↓nitrous oxide emissions) • Side-dress application to standing corn
July	✓	<ul style="list-style-type: none"> • Apply to just harvested forage fields (before regrowth; monitor rate) • Apply following wheat/spring cereal harvest (ideally with cover crops) • Apply to more distant fields; Sell manure to cash crop neighbours; or trade manure for straw, etc. • Apply following processing crops/edible beans/corn silage (ideally with cover crops)
August	✓	
September	<ul style="list-style-type: none"> X X 	
September	✓	<ul style="list-style-type: none"> • Pre-till tile-drained fields with large cracks/macropores (or apply < 4,000 gal/ac) • Avoid application on areas of concentrated flow and grassed waterways
October	✓	
November	<ul style="list-style-type: none"> X X 	
December	✓	<ul style="list-style-type: none"> • Apply manure only when immediate incorporation is possible • Avoid application on areas of concentrated flow and grassed waterways • Avoid application on frozen or snow covered fields • Where winter application is unavoidable (Contingency planning): <ul style="list-style-type: none"> o Access additional storage (neighbour, previous livestock facility) o Apply on fields protected from upslope runoff o Apply to fields with little slope (< 5 %) and no direct access to water sources o Keep liquid loading as low as possible (reduced rates) o Apply to fields with residue cover or living crops o Apply early in the winter when soil still has some warmth beneath the surface
January	X	
February		
March		

For copies of this article and a complete catalogue of previously published columns, check our website www.ccaontario.com under the "Communications" tab.

Christine Brown is a Field Crop Sustainability Specialist with OMAFRA. For any questions, she can be reached at christine.brown1@ontario.ca

There are over 650 Certified Crop Advisers (CCA) in Ontario. Each CCA has demonstrated their knowledge about Ontario crop production by passing the required exams. In addition, they have the crop advisory experience, the education, the commitment to continuing education and have signed a comprehensive code of ethics, which places the grower's interests first.



This industry driven program helps ensure that Ontario crop producers are well served by those providing their crop production advice. This article was written by one of those CCA's.