



Nitrogen Nutrition for Winter Wheat

by Dr. Tom Bruulsema, CCA-ON

The nitrogen nutrition of winter wheat is key to its yield and quality. Management choices for the right source, rate, time and place of nitrogen application support the sustainable intensification of wheat production needed to continue improving food and nutrition security for people.

Source: Since winter wheat often needs a topdress application, choosing the right source can help minimize ammonia loss. Urea on the soil surface has high potential to lose ammonia. The proportion lost increases with rate of application and soil pH, and can be influenced by soil moisture. Sources containing urea can be treated with a urease inhibitor—or protected with a slow-release coating—to slow hydrolysis, reduce loss, and supply more ammonium and less nitrate for plant uptake. As the levels of carbon dioxide in the air increase, wheat increasingly favors the uptake of ammonium over nitrate. Ammonium sulfate and ammonium nitrate sources are less prone to loss. The choice between fluid urea-ammonium nitrate and granular urea is tricky. Fluids can be applied more uniformly. Less ammonia is lost with fluid, but more leaf burn can occur. Leaf burn can be minimized with streamer nozzles and early application. Inhibitors or controlled-release forms do not always pay for themselves in terms of yield increase, but the benefits of reduced loss to the environment should also be considered.

Rate: The rate to apply depends on yield potential and previous crop. Rate predictions can also be aided using sensors that provide some measure of the canopy cover and its greenness. The optimum rate depends on cultivar as well. Hard red cultivars require more nitrogen for optimum protein levels. Cultivars susceptible to disease require lower rates. Production systems with weather-specific use of fungicide and plant growth regulators can produce higher yields with higher rates of applied nitrogen. For example, research conducted in Ontario, Canada from 2008 to 2010 demonstrated that, when combined with a fungicide strategy

that controlled disease, increasing rates of N in the spring topdress from 90 to 150 pounds per acre raised average yields from 90 to 112 bushels per acre in soft red winter wheat. These higher yields were accompanied by changes in grain quality considered desirable by millers, including increased protein and decreased levels of fungal toxins. Recent research in China and Turkey found that increasing rates of nitrogen increased the concentration and bioavailability of the micronutrient zinc in wheat grain. Considering the widespread distribution of zinc deficiencies in the human diet, increased zinc is a good thing.

Time: At wheat seeding in the fall, soils often contain sufficient nitrogen to start the crop. Exceptions occur, such as in sandy soils, or soils that have recently grown a nitrogen-depleting crop. Going into winter, the plant should not be deficient, but excess nitrogen can lead to disease infection and winterkill. Thus the decision to apply a small amount at seeding should be guided by a soil test. Topdress applications need to be timed according to the growth condition of the crop. Thin stands may benefit from early applications in the tillering phase, but only a low rate should be applied to prevent excessive tillering. By the beginning of stem elongation, Zadoks growth stage 30, most of the nitrogen should be applied, since from this stage the crop takes it up rapidly. A split application at both those stages may sometimes pay. After heading, nitrogen uptake slows down, but foliar applications, or controlled-release forms applied earlier, boost protein in the grain.

Place: the logical place for nitrogen fertilizer at seeding is with the seed. Wheat seedlings can tolerate the low rates required at planting. For topdress fertilizer, broadcast is often the only choice. Ensuring uniformity of broadcast is important for avoidance of uneven maturity. Uneven applications lose yield to both deficiency and lodging.

Nitrogen nutrition for wheat offers big opportunities to sustainably improve food and nutrition security.

For copies of this article and a complete catalogue of previously published columns, check our website www.ccaontario.com under the “Agronomic Articles” tab.



Dr. Tom Bruulsema is a Certified Crop Adviser employed by the International Plant Nutrition Institute as their Northeast Director located in Guelph. Contact Tom at (519) 835-2498 or tom.bruulsema@ipni.net

There are over 500 Certified Crop Advisers (CCA) in Ontario. Each CCA has demonstrated their knowledge about Ontario crop production by passing the required exams. In addition, they have the crop advisory experience, the education, the commitment to continuing education and have signed a comprehensive code of ethics, which places the grower's interests first.

This industry driven program helps ensure that Ontario crop producers are well served by those providing their crop production advice. This article was written by one of those CCA's.